Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.



RECEIVED

The table below shows the percentage of normal precipitation for the spring months. March to May combined, and for June, July, and August separately, for 1930, and for six other outstanding droughts in the United States. A comparison of the figures shows that the 1930 drought, when its length, area covered, and deficiency in precipitation are considered, to be the most severe of record. In the middle Atlantic area this drought began during the winter months.

In 1881 June was very dry in the Southwest and July and August in the Central Valleys and the East. In 1894 severe droughty conditions were confined to the Central Valleys and the Northwest, while in 1901 the Central Valleys, especially the western Corn Belt, suffered most. In 1911 the greatest deficiencies in precipitation occurred rather early in the season, while in 1916 and 1924 droughts covered comparatively small areas.

Mch. 1930 Mch. 1881 Mch. 1090 to 1911 to 1916 Mch. 1094 Mch. 1090 to 1911 to 1916 Mch. 1094 May June July Aug. May July Aug. M			Percentage of Normal	Precipitation
May June July Aug. May July Aug. Aug. Aug. Aug. Aug. Aug. Aug. Aug.	36-3-	: Mch : Mc		Mch. Mch.
May June July Rug. May June July Aug. May July Aug.			2000	
Md. 64 85 32 22 79 182 54 26 110 60 61 67428 79 123 135 64 105 61 230 98 151 125 151 137 107 77 98 124 61 74 38 45 68 74 74 27 72 46 89 76 138 108 104 198 80 83 68 136 79 135 121 73 137 107 77 98 115 77 121 137 107 77 98 115 77 121 137 107 77 98 115 77 131 137 107 77 98 115 77 121 137 107 77 98 115 145 130 105 115 145 130 105 115 115 137 107 77 98 115 115 130 105 115 115 130 105 115 115 115 115 115 115 115 115 11		May June July Aug. May	y June July Aug May Jur	ine July Aug May June July May May June July May May June July May May June July May May May May May May May May May Ma
Pa. 80 102 50 40 81 171 50 40 130 64 53 43139 86 89 162 77 115 64 181 100 174 118 26 129 114 57 88 N.J. 68 114 87 60 97 153 42 24 113 63 37 55148 44 29 196 80 146 79 178 92 114 118 26 129 114 57 88 95 N.Y. 97 116 77 47 86 97 62 40 105 93 67 43145 93 112 135 72 110 72 122 114 128 85 58 108 76 88 95	Md. 64 85 32 22 Va. 61 74 38 45 W.Va. 64 64 41 50 Ky. 51 45 30 50 Ohio 64 58 41 42 Ind. 63 68 53 34 Ill. 57 87 30 49 Mo. 59 79 24 66 Ark. 94 21 20 15 Tenn. 86 32 54 60 Ala. 77 47 74 75 Miss. 96 16 50 80 La. 79 12 60 70 Tex. 99 64 42 50 Okla. 95 97 32 90 Kans. 94 59 95 75 Nebr. 130 84 44 Iowa 78 129 39 50 S.Dak 90 74 37 90 N.Dak 81 89 42 68 Minn. 88 95 71 Wis. 83 136 70 25 Mich. 84 106 46 25 Pa. 80 102 50 N.J. 68 114 87	79 182 54 26 110 68 74 74 27 72 89 131 90 30 84 73 123 49 33 77 78 147 95 34 82 85 134 38 62 84 80 114 38 44 88 92 105 37 22 133 89 100 36 44 81 111 79 33 147 76 83 54 38 80 84 71 23 56 43 100 114 73 85 63 63 127 100 98 21 181 53 100 114 73 85 63 63 159 134 85 54 55 159 134 86 93 127 106 159 159 179 106 159 159 179 189 150 155 97 75 13 150 155 97 75 13 151 150 150 150 150 150 150 150 150 150	60 61 63128 79 46 89 76138 108 73 48 52144 126 68 49 81 80 100 65 41 55 64 116 68 41 55 66 85 68 41 55 66 85 70 43 33 105 124 79 33 70 55 93 15318 65 71 93 15318 60 72 79 81 200 76 39 73 48 41 25 86 138 74 84 41 25 86 138 75 69 18 39 77 42 83 77 114 116 83 33 90 46 18 83 77 114 116 83 33 90 46 18 83 83 64 53 43139 86 83 37 55148 41	123 135 64 105 61 230 98 151 125 51 153 150 57 95 8

^{*} Data for August, 1930, based on daily telegraphic reports of first order stations up to and including August 29, and subject to revision.

